

## Prevalence and Associated Factors of Postnatal Care Utilization among Women Attending Immunization Clinic in Health Centers in Hawassa City, Southern Ethiopia, 2019

Eleni Tesfaye Tegegne<sup>1</sup>,  
Mekibib Kassa Tessema<sup>2</sup>,  
Kaleab Tesfaye Tegegne<sup>3\*</sup>

- 1 College of Medicine and Health Science, School of Nursing, University of Gondar, Gondar, Ethiopia
- 2 Leishmania Research and Treatment Center, University of Gondar, Gondar, Ethiopia
- 3 Department of Public Health, Hawassa College of Health Science, Hawassa, Ethiopia

**\*Corresponding author:**

Kaleab Tesfaye Tegegne

✉ kaleabtesfaye35@gmail.com

Department of Public Health, Hawassa  
College of Health Science, Hawassa, Ethiopia

**Citation:** Tegegne ET, Tessema MK, Tegegne KT (2021) Prevalence and Associated Factors of Postnatal Care Utilization among Women Attending Immunization Clinic in Health Centers in Hawassa City, Southern Ethiopia, 2019. *J Rep Endo Infert.* Vol.6 No.4: 24.

### Abstract

**Introduction:** Postnatal care is a care provided to women and their babies within 42 days after delivery. A large proportion of maternal and neonatal deaths occur during the first 48 hours after delivery, and these first two days following delivery are critical for monitoring complications arising from the delivery. World Health Organization (WHO) recommends that after an uncomplicated vaginal birth in a health facility, healthy mothers newborns should receive care in the facility for at least 24 hours after birth. If birth is at home, the first postnatal contact should be as early as possible within 24 hours of birth.

**Objective:** To assess prevalence and associated factors of postnatal care utilization among women attending immunization clinics in health centers, Hawassa city, southern Ethiopia, 2019.

**Methodology:** Institution based cross sectional study design was done. Study was conducted in all health centers. The total sample sizes were 306 women attending immunization clinic at health centers. A random sampling technique was employed to select the study subjects. Data collection was conducted from study subjects using pretest questionnaires. The questionnaire was developed in English. Ethical clearance was obtained from the research unit of Rift Valley University. Data was entered and analyzed using SPSS version 20 for windows. Frequency distribution table and statically graph were used to describe the variables of interest. The associated factors of postnatal care utilization were identified using bivariate and multivariate analysis.

**Result:** Multi parous women who had PNC experience were three times more likely to utilize PNC service for the current delivery than those primi Para women [AOR=2.8, 95%CI (1.36-5.8)]. On the other hand, those multi parous women who hadn't PNC experience were almost 50% less likely to utilize PNC for the current delivery than those primi Para women [AOR=0.43, 95%CI (0.20-0.92)]. Finally, those women who were discharged from the institution after delivery at less than 6 hrs of stay were almost 80% less likely to utilize PNC than those women who stayed more than 24hrs [AOR= 0.22, 95%CI (0.06-0.83)].

**Conclusion:** The overall prevalence of PNC service utilization in this study was relatively good as compared to the HSDP IV report for Hawassa city in the 2006 EFY. PNC counseling and provision of appointment, counseling on danger sign, past experience of PNC utilization, and less than 6 hours stay at health institution before discharge were found to be statistically significant for the current PNC service Utilization,

**Keywords:** Ethiopia; Immunization; Postnatal care; Women

**Received:** June 08, 2021, **Accepted:** July 22, 2021, **Published:** July 28, 2021

### Introduction

#### Background information

Postnatal care is a care provided to women and their babies within 42 days after delivery [1]. A large proportion of maternal

and neonatal deaths occur during the first 48 hours after delivery, and these first two days following delivery are critical for monitoring complications arising from the delivery [2]. World Health Organization (WHO) recommends that after an uncomplicated vaginal birth in a health facility, healthy mother's

newborns should receive care in the facility for at least 24 hours after birth. If birth is at home, the first postnatal contact should be as early as possible within 24 hours of birth.

At least three additional postnatal contacts are recommended for all mothers and newborns, on day 3 (48- 72 hours), between days 7-14 after birth, and six weeks after birth [1].

Whereas Federal Ministry of Health (FMoH) of Ethiopia recommends three PNC care visits at 6 hour, 24 hours, 3 days, 6 days and 6 weeks [3,4]. The health of mothers is mostly regarded as an indicator the health of the society [2]. Postnatal care is regarded as one of the most important maternal healthcare services for the prevention of impairments and disabilities resulting from childbirth [5].

Lack of care during puerperium may result in death or disability as well as missed opportunities to promote healthy behaviors, affecting women, newborns, and children [6]. Millions of mothers more suffer severe illness each year, and unknown numbers are affected with lifelong disabilities. Some of the long-term maternal complications in the postnatal period include chronic pain, impaired mobility, damage to the reproductive system and infertility [7]. Some women suffer genital prolapses after bearing several children. This condition is extremely uncomfortable and can lead to other complications in future pregnancies if not properly addressed in the postnatal period [5].

Ethiopia is one of the countries with high maternal mortality. The MMR was 871 per 100,000 in the year 2000; it was 673 per 100,000 live births in 2005 and 676 per 100,000 in 2011. Maternal deaths represent 30% of all deaths to women age 15-49, compared with 21% in the 2005 EDHS and 25% in the 2000 EDHS [6]. Poor women in remote areas are the least likely to receive adequate health care. This is especially true for regions with low numbers of skilled health workers, such as sub-Saharan Africa and South Asia.

During the past decade, only 46% of women in low-income countries benefit from skilled care during childbirth. This means that millions of births are not assisted by a midwife, a doctor or a trained nurse. Other factors that prevent women from receiving or seeking care during pregnancy and childbirth are: poverty, distance, lack of information, inadequate services, and cultural practices.

The level of postnatal care coverage is extremely low in Ethiopia 18 percent. Of this only 13 percent of women received postnatal care within two days, as recommended. Among women who received a postnatal checkup, 8 percent were examined within 4 hours of delivery, 3 percent within 4-23 hours, 2 percent within 1-2 days, and 5 percent within 3-41 days of delivery [7]. Low coverage of postnatal care in Ethiopia is causing to continuous high maternal and new born morbidity and mortality that affects MDGs 4 and 5 [8]. It is also challenge for planning and implementing of PNC as well as many opportunities are missed low PNC coverage including exclusive breastfeeding, PMTCT, providing of family planning and maternal and new born care [9].

The burden of low of postnatal care in Ethiopia is even greater for the large rural population due to poor access and utilization of maternal and new-born health services [10].

## Statement of the problem

The World Health Organization (WHO) stated that Postnatal care (PNC) is defined as a care given to the mother and her newborn baby immediately after the birth of the placenta and for the first six weeks of life [5]. It is important because during this period the reproductive organs return to their pre gravid state. Lactation is started, the mother recuperates from the physical and emotional stress of Labor and the family adjusts to the new baby.

PNC services are to support the mother and her family in the transition to a new family consultation, prevent, early diagnose and treat complications of the mother and newborn, refer the mother and new born for specialist care when necessary, counsel on baby care, support breastfeeding, maternal nutrition, provide contraception service, and immunize the infant. With limited resources, contact with the health care system at least during the first twenty four hours and before the end of the first week would be the most effective strategy [11].

This period is generally the most neglected in developing countries mothers and new born babies do not receive postnatal care services from a skilled health care provider during the first few days after delivery [12,8]. In developed countries virtually all women and their infants receive PNC, even though the nature and frequency of this care varies considerably [8]. However, in developing countries even the need for care and support after birth was less recognized and approximately one-third of women in sub-Saharan Africa give birth in facilities, and no more than 13 percent receive PNC within two days of delivery. Whether women deliver at home or in a facility, PNC services are often absent. Moreover, PNC services, where available, often lack essential elements of care required for the optimum health of the mother and her newborn [8,7].

To assess the extent of postnatal care utilization, the 2011 EDHS asked respondents whether they had received a health checkup after the delivery, the timing of the first check, and the type of health provider for their last birth in the two years preceding the survey, and it was found that the level of postnatal care coverage was extremely low in Ethiopia. The great majority of women (92 percent) with a live birth in the preceding five years did not receive a postnatal checkup. Among women who received a postnatal checkup, 4 percent were examined within 4 hours of delivery, 2 percent within 4-23 hours, 1 percent within 1-2 days, and 2 percent within 3-41 days of delivery. In total, only 7 percent of women received postnatal care within two days, as recommended while half (51.5%) of them in Addis Ababa had ever used postnatal care [3].

Despite the fact that no previously done studies existed in the study area (Hawassa, SNNPR, Ethiopia) on factors associated with PNC service, 2014 report indicated that the level of PNC coverage was extremely low in SNNPR (15.4%) and only few (11.1%) of mothers were received PNC service within 48 hours after gave birth (Ethiopia Mini Demographic and Health Survey .2014).

Therefore this study will be conducted for the assessment of prevalence of postnatal service utilization and associated factors in Hawassa city.

## Significance of the study

Even though PNC service utilization plays a critical role in reducing maternal and new born child mortality, little is known about its factors so, this study is planned to identify the possible associated factors for PNC service utilization it is important to make additional studies to describe the magnitude and associated factors of low level of postnatal care utilization.

The study will be conducted for the assessment of prevalence of postnatal service utilization and associated factors in health centers in Hawassa city and will be used as baseline data for further research studies

The factors associated with utilization of PNC services still need strong due attention to be researcher so as to improve its utilization and also it is convincing that the available knowledge about the service utilization is insufficient. This study therefore intends to assess these anticipated bottlenecks of PNC service utilization and in addition fill some of the gaps of previously carried out literatures on similar topic with the view of improving utilization of the services.

## Methods and Materials

### Study area and period

The study was conducted at all health centers in Hawassa city administration. Hawassa city, which is 273 km away from Addis Ababa in the south The Health centers, offers health services including maternal and child health services. Some of these are outpatient, TB/ leprosy, antenatal care, delivery, postnatal care, and family planning services (Hawassa city health office, unpublished). We estimate the number of PNC service users for the previous three consecutive months in 10 health centers of Hawassa city which was 1000(Hawassa city health office, unpublished). The study was conducted from June -July 2019.

### Study design

Institutional based cross sectional study was conducted.

### Source and study population

**Source Population:** All women who deliver a baby and attend maternal and child health clinics within six week of delivery in Hawassa city.

**Study Population:** All selected women, who delivered a baby and come to maternal and child health clinics at the ten selected health centers in Hawassa city and fulfill the inclusion criteria.

### Inclusion and exclusion criteria

**Inclusion Criteria:** Women who visited the selected health centers at maternal and child health clinics during postpartum period and gave informed consent.

**Exclusion Criteria:** Women who had difficulty in communication due to severe mental disorder (problem)

### Sample size determination and sampling procedure

**Sample Size Determination:** Single population proportion

formula was used to calculate a sample size, by using 70% expected magnitude of postnatal care utilization (Regassa N, 2011). 10% Non-response rate were considered.

$$N = z^2 pq / d^2$$

Where  $z = 1.96$  (at 95% confidence interval)

$P = 51.4\%$  (magnitude of postnatal care utilization)

$$q = 1 - 0.514 = 0.486$$

$d = 0.05$  degree of accuracy or margin of error

$$\text{Hence } N_i = z^2 pq / d^2$$

$$N_i = (1.96)^2 (0.514) (0.486) / (0.05)^2 = 384$$

Since population is  $< 10,000$  we use correction formula

$$N = 384 / (1 + 384 / 1000) = 278$$

$$N_f = 278 + 10\% (278) = 306$$

Thus by adding 10% none response rate, the final sample size will be 306.

### Sampling technique

The study was conducted in all health centers of Hawassa city to represent all the inhabitants of the city. The total sample size was proportionally allocated for the 10 health centers based on the six month report of maternal and child health clinics client flow of each health center. All eligible women in each health centers were invited to participate consecutively during the study period till the required sample size achieved.

### Study Variables

**Independent variables:** Socio demographic characters: Age, Religion, Ethnicity, Current marital status, Monthly income, Occupation of the partner.

Maternal reproductive and obstetric factors: Gravida, Para, abortion, Number of alive children, Nature of pregnancy (planned Vs. Unplanned, supported vs. unsupported) Distance of health institution Mode of delivery, Place of delivery, ANC utilization, previous postnatal care utilization, length of stay in the facility after delivery.

**Dependent variable:** Post natal care utilization

### Data collection tools

Data was collected using structured Amharic interview questionnaire having three parts, the first part containing socio demographic characteristics. The second part of the questionnaire was assessing the reproductive and obstetrics characteristics of the mother. The final part of the questionnaire contains asking prevalence of postnatal care utilization

### Data quality control

In order to maintain quality of the data, data collectors and supervisors was trained in data collection procedures. The questionnaire was carefully designed and prepared in English language first and then translated in to Amharic by language experts and again the Amharic version translated back to English

to make it consistent.

Finally, Amharic version was used to collect data. Before actual data collection time the questionnaire (tool) was pretested for validity and reliability on 5% of total sample size at yirgalem health center, there by possible adjustment or modification was made on the tool. The collected data was then reviewed and checked for completeness and consistency by the graduate students on a daily basis.

### Data processing and analysis

The data was checked, cleaned, coded and entered by using Statistical Package for Social Science (SPSS) version 20 statistical software for analysis.

Descriptive statistics was computed to determine the proportion of postnatal care service utilization. chi-square test was computed to assess statistical association between the outcome variable and independent variables using Odds Ratio; significant of statistical association were assured or tested using 95% confidence interval (CI) and p value (<0.05).

### Operational definitions

**Postnatal Care Utilization:** women had at least one check up by the health professional after delivery within 6 weeks of postpartum coming to the health facility. (Normally women are expected to visit the PN clinic during the first 24 hrs., 3rd day, 7th day, 14th day and 42nd days).

**Postnatal Care:** refers to the assistance or care given to mother and baby by health workers during the postnatal follow up period such as counsel on baby care, support breastfeeding, maternal nutrition, provide contraception service, and immunize the infant.

### Ethical Considerations

Permission letter was written by Rift Valley University, department of nursing to health centers. A formal letter, from Rift Valley University was submitted to the head of health centers. Women who attend immunization clinic were informed and requested their permission to conduct the study. The purpose of study was explained to the study subject at the time of data collection and verbal consent was taken from participants. Information obtained was kept confidential

## Results

### Socio demographic characteristics of the participants

A total of 306 women participated in the study with a response rate of 100%. Almost half, 187 (61.1%) of the Participants were found between the age of 25 to 29 years. The mean age of the participants was 26.7 years (+SD=4.4). 272 (88.89%) of them were orthodox Christian followers and 294 (96.07%) of them were married. With respect to level of education only 79 (25.81%) of the respondents had college and above education. The average monthly house hold income was 3800 ETB. More than half 242 (79.08%) of the participants were using taxi for manse of transportation to the health facilities (**Table 1**).

**Table 1:** Socio-demographic characteristics of the Women of Interviewed in Hawassa Health Centers 2019, (n=306).

Variables	Frequency (no)
<b>Age</b>	
<20 years	18
20-24 years	110
25-29 years	87
30-34 years	81
>=35 years	26
<b>Religion</b>	
Orthodox Christian	172
Muslim	92
Protestant Christian	52
Others	6
<b>Ethnicity</b>	
Sidama	108
Oromo	46
Tigre	36
Gurage	91
Others	41
<b>Marital status</b>	
Married	194
Single	57
Divorced	87
<b>Educational status</b>	
No formal education	65
Primary school	98
Secondary school	64
College and above	79
<b>Paternal educational status</b>	
No formal education	23
Primary school	66
Secondary school	109
College and above	108
<b>Occupation of participants</b>	
House wife (unemployed)	163
Employed	143
<b>Family monthly income**</b>	
<500 Eth Birr	41
500-999 Eth Birr	21
1000-1499 Eth Birr	38
1500-1999 Eth birr	40
>=2000 birr	166

### Reproductive characteristics of the participants

More than 243 (79.41%) of the participants didn't have history of abortion. Half 207 (67.64%) of the participants were primigravida and a quarter 201 (65.6%) of the current pregnancy were unplanned. All most all 212 (69.28%) respondents had ANC follow up at least one, on top of that more than half 223 (72.87%) had four times and above follow up as show below in **Table 2**.

### Labor, delivery and postpartum characteristics of the participants

Institutional delivery among the study participants were 202

**Table 2:** Reproductive characteristics of the Women of Interviewed in Hawassa Health Centers.

Variables	Frequency( no)
<b>History of abortion</b>	
Yes	79
No	243
<b>Parity</b>	
Para I	207
Para 2-4	92
>=5	7
<b>Nature of index pregnancy</b>	
Planned and supported	131
Unplanned but supported	120
Unplanned and unsupported	55
<b>History of ANC Yes</b>	
Yes	252
No	56
<b>Number of ANC visits</b>	
One times	8
Two times	25
Three times	154
Four times	186
Five and above times	37
None	12
<b>Gestational age of index pregnancy</b>	
Term	159
Preterm	10
Post term	53
<b>Number of pregnancy</b>	
Singleton	105
Twin	17
<b>Sex of the neonate</b>	
Male	206
Female	100
<b>Types of feeding</b>	
Exclusive breast feeding	275
Formula feeding	47
<b>APH during pregnancy</b>	
Yes	12
No	210
<b>Hypertensive disorders during pregnancy</b>	
Yes	6
No	300
<b>PROM and choriarnioties during pregnancy</b>	
Yes	17
No	299

(66.01%) and vaginal delivery accounts more than three fourth 245(80.06%) of the participants. The mean duration of institution stay before discharge after delivery was 26.7 hrs. (+SD=41hrs.),

however the mode is 6 hrs. Nearly half 192(62.74%) of women stay 6 to 11hrs before discharge. Three fourth 221(72.22%) of the participants were given appointment for postnatal care by the health care professionals before discharge. The remaining 101(33%) participants were not informed to have postnatal care by the health professionals before discharge from the health institution. Among women who gave birth in the health institution, and 258 (84.31%) of them were advised about any danger signs of postpartum period before discharge (**Table 3**).

**Table 3:** Labor, delivery and postpartum characteristics of the Women of Interviewed in Hawassa Health Centers o Hawassa city, 2019 (n=306)

<b>Place of delivery</b>	
Home	16
Health center	120
Government hospital	123
Private hospital	51
Private clinic	12
<b>Delivery attended by</b>	
Midwife(Nurse)	148
Physician	60
Traditional birth attendant/family	14
<b>Mode of delivery</b>	
Vaginal delivery	245
Instrumental delivery	10
Cesearn section	67
<b>Episiotomy/perianal tear during vaginal delivery (N=306)</b>	
Yes	202
No	104
<b>Types of caesarean section(N=67)</b>	
Emergency	42
Elective	26
<b>Duration of institution stay after delivery*</b>	
< 6hrs	21
6-11 hrs	102
12-23hrs	73
>=24 hrs	108
<b>Did you given appointment for postnatal care before discharge</b>	
Yes	205
No	101
<b>Appointed within 48 hrs of discharge(N=306)</b>	
Yes	40
No	266
<b>Appointment after one week</b>	
Yes	147
No	74
<b>Appointed within 2-4 weeks</b>	
Yes	17
No	205

## Prevalence and characteristics of postnatal care utilization

The most frequent place of postnatal care were at health centers 199(65.03%) followed by the government hospitals 51(16.66%). The proportion of postnatal care visit across 48hrs of discharge, after one week of discharge and at six weeks of postpartum were 28(0.91%), 199(65.03%) and 143(46.73%) respectively. With regard to the frequency of postnatal care visit, 159 (51.96%) of participants had visited once, 112(36.60%) women had two time, and the remaining 6(0.19%) were having three or more (**Table 4**).

**Table 4:** Prevalence and characteristics of postnatal care utilization of interviewed women in selected government Health Centers of Hawassa city, Ethiopia, 2019, (n=306).

Variables	Frequency
<b>Did you have postnatal care N=306</b>	
Yes	161
No	145
<b>Did you have postnatal for the previous delivery, if you gave birth N=224</b>	
Yes	147
No	77
<b>Place of postnatal care N=277</b>	
Health center	199
Gov't hospital	51
Private hospital	23
Private clinic	4
<b>Postnatal care within 48hrs of discharge N=277</b>	
Yes	28
No	149
<b>Postnatal care after one weeks of discharge N=277</b>	
Yes	199
No	78
<b>Postnatal care at six weeks of postpartum N=277</b>	
Yes	143
No	134
<b>Number of postnatal care visit N=277</b>	
One times	159
Two times	112
Three times	6
<b>Did your baby was with you during postnatal care N=277</b>	
Yes	262
No	15
<b>Contraceptive use at 14th weeks</b>	
Yes	255
No	167
<b>Contraceptive initiation period *</b>	
Before 6 weeks (42 days)	23
At 6 weeks (42 days)	211
After 6 weeks	21

Concerning what was done for the women during postnatal care visit we found that nearly half 53.6% of the respondents provided contraceptives, 35.4% had physical examination, 22.4% advice on danger signs, 10.5% provided TT vaccination, and 5.8% had laboratory investigations.

## Associated factors of postnatal care utilization

During bivariate logistic regression analysis, those variables who had significant association to the dependent

Variables with P-values of less than 0.2 were entered to multivariate logistic regression. Out of twenty five independent variables categorized under Socio demographic, reproductive and obstetrics, labor and postpartum characteristics, eight variables namely educational status, nature of pregnancy, ANC, place of delivery, whether or not PNC appointment given, whether or not advise on danger signs given by the health care provider, previous history of PNC, and duration of institution stay before discharge had showed association with p-values of less than 0.2. After we entered these 8 variables in to the logistic regression the following four variables provision of PNC appointment, counseling on danger signs by the health care provider, previous history of PNC, and duration of institution stay before discharge had showed statistical significant association with PNC utilization.

Accordingly, those women who were counseled and given appointment for postnatal care utilization had over thirty two times [(AOR=32.6, 95%CI (14.7-72.3)] more likely to utilize postnatal care than those women who didn't informed by the health care providers before discharge. The odds of having postnatal care visit for those women who were counseled about any of danger signs by the health care providers before discharge were two times more likely to have PNC visit than their counterparts [AOR (1.95, 95%CI (1.05-3.64)]. Similarly those multiparous women who had PNC experience were three times more likely to utilize PNC service for the current delivery than those primipara women [AOR=2.8, 95%CI (1.36-5.8)]. On the other hand, those multiparous women who hadn't PNC experience were almost 50% less likely to utilize PNC for the current delivery than those primipara women [AOR=0.43, 95%CI (0.20-0.92)]. Finally, those women who were discharged from the institution after delivery at less than 6 hrs of stay were almost 80% less likely to utilize PNC than those women who stayed more than 24hrs [AOR= 0.22, 95%CI (0.06-0.83)] (**Table 5**).

## Discussion

This study tried to assess the prevalence and associated factors of postnatal care utilization among women who delivered a baby and attending immunization on the 10<sup>th</sup> and 14<sup>th</sup> weeks of postpartum at five selected health centers in Hawassa city. Most of the participants in this study were multiparous 208(49.3%), illiterate 65(15.4%), and 394(93.4%) were married.

This study indicated that two thirds (65.6%) of the participants had received a health checkup after delivery despite the fact that almost all women were given birth at health institution 406(96.2%). But this figure is higher when compared with the 2011 EDHS postnatal care utilization rate of 51.5% and 82.3% of institutional delivery rate respectively [12]. This improvement

**Table 5:** Logistic regression analysis of factors associated postnatal care utilization of the Women of Interviewed in Health Centers of Hawassa city , Ethiopia, 2019 (n=306).

Variable	PNC UTILIZATION			
	YES	NO	COR(CI)	AOR(CI)
No formal education	38	27	0.77(0.39-1.52)	1.56(0.57-4.4)
Primary school	99	59	0.92(0.53-1.62)	1.88(0.8-4.45)
Secondary school	89	31	1.56(0.85-2.9)	2.22(0.92-5.36)
Above secondary school	51	28	1	1
<b>Nature of pregnancy</b>				
Planned and supported	113	88	2.61(0.78-8.75)	1.0(0.17-5.91)
Unplanned and supported	59	41	1.73(0.49-6.04)	0.6(0.01-3.61)
Unplanned and unsupported	5	6	1	1
<b>ANC</b>				
Yes	173	99	2.95(0.82-10.6)	5.93(1.00-30.5)
No	4	6	1	1
<b>Place of delivery</b>				
Home	2	14	<b>0.07(0.011-481)</b>	0.31(0.01-7.63)
Health center	68	52	1.62(0.47-5.58)	1.01(0.16-7.5)
Gov't hospital	66	57	.58(0.17-2.02)	0.59(0.09-4.1)
Private hospital	33	18	.92(0.24-3.47)	0.74(0.10-5.43)
Private clinic	8	4	1	1
<b>Appointment given for PNC</b>				
Yes	166	55	<b>39.57(19.85-78.8)</b>	<b>32.6(14.7-72.3)**</b>
No	11	90	1	1
<b>Women counseled any danger sign on discharge</b>				
Yes	102	56	<b>4.28(2.79-6.56)</b>	<b>1.95(1.05-3.64)**</b>
No	75	89	1	1
<b>PNC experience in the previous delivery</b>				
Multi Para Yes	123	24	<b>3.13(1.85-5.27)</b>	<b>2.8(1.36-5.8)**</b>
Multi Para No	31	46	<b>0.41(0.24-0.70)</b>	<b>0.43(0.20-0.92)**</b>
Primi Para	83	75	1	1
<b>Duration of institution stay before discharge</b>				
<6 hrs	8	13	<b>0.3(.12-.78)</b>	<b>0.22(0.06-0.83)**</b>
6-11 hrs	84	58	1.13(.69-1.83)	0.81(0.36-1.83)
12-23hrs	48	25	0.94(.51-1.72)	0.34(0.13-0.88)
>=24hrs	86	42	1	1

might be due to the time difference and the presence of diverse intervention to improvement in accessing and utilizing maternal health care service. This study finding on PNC prevalence was also higher than three similar studies conducted in Enderta District, Tigray,49.7%(24), Jabitena District, Amhara Region (20.2%)and southern Ethiopia (37.2%). The discrepancies might be associated with the variation in the study areas and study subjects.

Similarly the postnatal care utilization of this study finding also a bit higher when compared to other studies conducted in African and Asian countries ; the prevalence of not using postnatal care were Bangladesh (73%), Nepal (72%), Rwanda (71%), Burkina Faso (44%), Cambodia (46%), Haiti (55%), and Kenya (46%) Malawi (41%), Mali (49%), Nigeria (46.5%), Uganda (57%) and

Zambia (41%) [13]. However the findings were lower when compared to a study conducted in Brazil (77%) and many other developed countries [6,14].

Concerning the quality of postnatal care provided by the health professional, we found that about half (53.6%) of the women we provided only contraceptives and less than one in three women received the remaining postnatal care services such as counseling on danger signs, physical examination, and health promotion. similarly, the postnatal care provided to their babies mainly focused on providing immunization (98.1%). The remaining services accounts for less than 20%. This is also true in a study done in southern Ethiopia.

The key associated factors for PNC utilization in our study were PNC appointment given by the health provider, counseling of

women on danger signs, history of previous PNC utilization and length of health institution stay before discharge. Women who were counseled and given appointment for postnatal care service were utilized the PNC service over thirty two times [(AOR=32.6, 95%CI(14.7-72.3)] higher than to those women who didn't informed about the PNC service on discharge. This is also supported by the reason given by those women who didn't utilized PNC during this study. Similarly, a study done in northern Ethiopia found that those women who had got information about postnatal care services from HEW and Midwife/Nurse were 24.87

And 37 times more likely to attend postnatal care service respectively compared to those women's who had got information from other sources. This finding may leads to a conclusion that the PNC service utilization is strongly influenced by the knowledge of women on postnatal care benefits.

The other major factors predicting postnatal care service utilization was counseling of women about danger sign during the PN period. Those women who were informed about any of the danger signs that occur during the PNC period were two times more likely to utilize postnatal care than their counterparts [AOR (1.95,95%CI (1.05-3.64)]. This finding was in line with a study conducted in Amhara Region. Mothers who were knowledgeable at least for one postpartum obstetric danger sign were more likely to utilize PNC service as compared to those who did not. Similar findings also reported from studies conducted in Nepal, and Uganda. This can be explained by the fact that awareness of obstetric danger signs is an important factor in motivating women and their families to attend health care service at the earliest opportunity with the intention of prevention, early detection and getting managed their obstetric danger signs.

The other major factors predicting postnatal care service utilization were previous history of postnatal care utilization and duration of institution stay before discharge. Those women who were discharged from the institution after delivery with less than 6 hrs. Of stay were almost 80%less likely to have PNC than those women who stayed more than 24 hrs. [AOR= 0.22, 95%CI (0.06-0.83)]. Those women who had gave birth before and had PNC were three times more likely to have PNC currently than those women who gave birth only once [ AOR=2.8, 95%CI (1.36-5.8)].

This strong positive association of PNC services utilization with previous history can be attributed to the fact that women who had PNC in health institution have greater opportunity to get exposed to health education related to PNC services at the time of visit and thus get access to learn about the types, benefits and availabilities of PNC services during their stay in the health institutions.

Finally, unlike previous studies in Ethiopia which finds that educational status of the participants, having ANC and socioeconomic status of the participants were the determinants of PNC utilization, these factors and others were not the determinant factors of PNC utilization during this study. This could be mainly because the study conducted in the most urban city of the country and almost all women have similar access to information regarding PNC service through media or during their ANC follow up visits.

## Limitation of the Study

Study design was cross sectional so that cause and effect relationship of variables were difficult to ascertain.

Use of health professionals as data collectors may create bias as they might direct the respondents during the data collection.

## Conclusion

The overall prevalence of PNC service utilization in this study was relatively good as compared to the HSDP IV report for Hawassa city in the 2006 EFY. PNC counseling and provision of appointment, counseling on danger sign, past experience of PNC utilization, and less than 6 hours stay at health institution before discharge were found to statistically significant for the current PNC service Utilization.

## Recommendations

**Health Professionals:** Should be counseled about the danger signs of postnatal period and provided appointment. Should be stayed at least for 24 hrs. following delivery. Should give PNC appointment and counsel about danger sign during postnatal period.

**Health facility:** Should assure comprehensive and quality of postnatal care services. Should give continuous training for health care providers on PNC appointment and counseling about danger sign during postnatal period.

**Bureau:** Should emphasize the improvement of quality of service PNC utilization. Should prepare guide lines on danger sign during Post natal

## Authors Contributions

Kaleab Tesfaye Tegegne, Eleni Tesfaye Tegegne and Mekibib Kassa Tessema Was responsible for conceptualization, project administration, software, supervision, and development of the original drafting of the manuscript.

Kaleab Tesfaye Tegegne, Eleni Tesfaye Tegegne and Mekibib Kassa Tessema were participated in methodology, analysis and manuscript preparation.

All authors contributed with data analysis, critically revised the paper, and agreed to be accountable for their contribution.

## Acknowledgements

We would like to thank all the primary authors of the included articles

## Funding Sources

There is no source of funding

## Conflict of Interest

The author(s) declares no conflict of interest

## References

- 1 World Health Organization recommendations on Postnatal care of the mother and newborn, USA (2013).
- 2 Ethiopia Demographic and Health Survey (2011) Central Statistical Agency Addis Ababa, Ethiopia ICF International Calverton, Maryland, USA.
- 3 Ethiopia Mini Demographic and Health Survey (2014) Central Statistical Agency Addis Ababa.
- 4 Federal Democratic Republic of Ethiopia Ministry of Health (2010) Management protocol on selected obstetrics topics
- 5 Fikirte T, Walelegn W, Fekadu M (2011) Knowledge, Perception and Utilization of Postnatal Care of Mothers in Gondar Zuria District, Ethiopia: A Cross- Sectional Study. *Maternal Child Health J* 18: 2341-51.
- 6 Regassa N (2011) Antenatal and postnatal care service utilization in Southern Ethiopia: A population-based study. *Afr Health Sci* 11: 390-397.
- 7 Vishnu K, Mandira A, Rajendra K (2014) Factors associated with the utilization of postnatal care services among the mothers of Nepal: analysis of Nepal Demographic and Health Survey 2011. *BMC Women's Health* 14: 19.
- 8 Stella B, Adesegun F (2005) Determinants of use of maternal health services in Nigeria-looking beyond individual and household factors. *BMC Pregnancy and Childbirth* 9: 43.
- 9 Asma Y, Sarosh I, Riffat M (2013) Determinants of Postnatal Care Services Utilization in Pakistan- Insights from Pakistan Demographic and Health Survey (PDHS) 2006-07. *Middle East J Sci Res* 18: 1440-1447.
- 10 Berhe H, Tilahun W, Aregay A (2012) Utilization and associated factors of postnatal care in Adwa town, Tigray, Ethiopia: A cross sectional study.
- 11 Hailemariam A, Huruy A, Yohannes A (2013) Prevalence and Factors Associated with Post Natal Care Utilization in Abi-Adi Town, Tigray, Ethiopia: A Cross Sectional Study. *IJPBSF* 8: 1.
- 12 Desalegn M, Daniel B (2013) Birth preparedness and complication readiness among women of child bearing age group in Goba woreda, Oromia region, Ethiopia. *BMC Pregnancy and Childbirth* 14: 282.
- 13 USAID Maternal and Child health Program (2012) Cultural Barriers to Seeking Maternal Health Care in Ethiopia: A Review of the Literature.
- 14 Solomon T, Danika B, Abebe G (2014) Improving Coverage of Postnatal Care in Rural Ethiopia Using: A Community-based, Collaborative Quality Improvement Approach. *J Midwifery Womens Health* 2014 59: S55-S64.